

EWM Management

No EWM

EWM Present



Monitoring

Spring and fall monitoring of the littoral zone (consultant) and monitoring throughout the summer (volunteers)

Type of Infestation

(1) Isolated Plants
(RF = V-1)



(2) Small Beds
(RF = 1-2)



(3) Thick Beds
(RF = 2-3)



of Plants

(a) 0-25 (b) 25-50 (c) >50

Coverage Area

(i) Pre - 0.01 ac. (ii) 0.01 - 0.1 ac. (iii) 0.1 - 0.2 ac. (iv) 0.2 - 0.5 ac.

(X) Shallow <5-ft

(Y) Deep >5-ft

Physical
(hand, rake, snorkel)

(1,2,a,b,i,ii,X)

Can be performed at any time by volunteers, especially around docks and swim areas

Scuba/DASH
(diver)

(1,2,b,c,ii,iii,iv,X,Y)

To be done by trained professionals with volunteer and consultant support

Herbicide Application

(2,3,v,vi,X,Y)

(v) 0.2 - 3.0 ac. (vi) >3.0 ac.

To be applied by certified contractors before June 30th in areas delineated from previous fall bed mapping and adjusted as needed from spring bed mapping

ProcellaCOR
2,4-D/Triclopyr